

Rochester Gastroenterology Associates, LLP

Prasad Penmetsa, M.D., M.R.C.P.

Surinder Devgun, M.D.

Ari Chodos, M.D.

Amy Hayes, F.N.P.-C.

Sarah Pratt, F.N.P.-C.

www.rochestergastro.com

INSTRUCTIONS FOR COLONOSCOPY PREP WITH MIRALAX, DULCOLAX & GATORADE

Name _____ Procedure _____

YOU HAVE BEEN SCHEDULED FOR THE ABOVE EXAMINATION BY DR. _____
PLEASE REPORT TO:

Unity Hospital

1555 Long Pond Road
Rochester, NY 14626
Tel: (585) 723-7119

Rochester Endoscopy Center

20 Hagen Dr., Suite 330
Rochester, NY 14625
Tel: (585) 267-4040

Wilson Health Center

800 Carter Street
Rochester, NY 14621
Tel: (585) 922-0825

Rochester General Hospital

1425 Portland Avenue
Rochester, NY 14621
Tel: (585) 922-2990

REPORT AND REGISTRATION TIME:

DATE/DAY:

PLEASE REVIEW ALL INSTRUCTIONS UPON RECEIPT

You will need to obtain one (1) **238 Gram** bottle of MiraLax (Glycolax), one (1) small package of Dulcolax Laxative **tablets** (4 tablets only) and two (2) **32 oz** bottles of Gatorade, (**NO** Red or pink) Most people prefer 2 different flavors – as an alternative you may use **64 ounces** of any other clear liquid (apple juice, crystal light, lemonade, iced tea etc.)

MiraLax and Dulcolax are purchased over the counter.

Transportation arrangements must be made for someone to pick you up and accompany you home. **DUE TO THIS BEING A NEW YORK STATE LAW. NO EXCEPTIONS CAN BE MADE!**

You must arrange for a neighbor, friend, or family member to stay with you after your arrival home since sedation effects can last well into the evening.

If you are unable to keep your scheduled appointment, cancellation is appreciated **48 hours** in advance by calling (585)267-4040 or (585)227-1080 between **8:30 a.m.** and **4:30 p.m.** If you fail to notify us you may be charged. Please call these numbers if you have any further questions.

If you are taking *any* kind of blood-thinning medications, such as **COUMADIN, PLAVIX, or XARELTO** please inform our office at (585)267-4040 or (585)227-1080, **IMMEDIATELY**. ***These medications MAY need to be stopped as DIRECTED BY YOUR MEDICAL PROVIDER.***

Page 2 In order to assure optimal colonoscopy examination, the following guidelines are recommended:

1. **Stop** any extra dietary fiber, including Metamucil, Citrucel, Fibercon, or similar products for **1 week** before the procedure.
2. **Stop** all iron medications and vitamins containing iron for **1week** before the procedure.
3. **Do not eat** nuts, raisins, or foods containing seeds (strawberries, sesame, poppy, etc.) or uncooked vegetables **3 days** before the procedure.
4. **Take your regular medications on the morning of the examination, except insulin or other diabetic medications.** Please check with our office prior to taking any **INSULIN** or other diabetic medication on the morning of the examination.

DAY BEFORE EXAMINATION

1. You may have a *light* breakfast BEFORE 8 am. Begin a clear liquid diet AFTER 8 am. **NO solid food is allowed.** Clear liquids include water, strained fruit juice without pulp (apple, white grape, lemonade), clear broth or bouillon, coffee and tea (without cream or non-dairy creamer), Gatorade (**NO** red or pink), carbonated and non-carbonated soft drinks, Kool-Aid (**NO** red or pink), Jell-O (**NO** red or pink), and popsicles (**NO** red or pink).
2. At **1 pm** take only **2 Dulcolax tablets** with water and continue on clear liquids. Then add **1/2** of the MiraLax powder to one **32 oz** bottle of Gatorade (**NO** red or pink) and mix until dissolved. Repeat this step with the other **1/2** of MiraLax powder and second **32 oz** bottle of Gatorade, then chill if desired.
3. At **2 pm** begin drinking the MiraLax/Gatorade solution. Drink this solution slowly, as tolerated, completing it by **7 pm**. It may take several hours for results. *It does not work very quickly*
4. At **5 pm** take the **2 additional Dulcolax tablets, and continue drinking clear liquids, throughout evening. THIS IS VERY IMPORTANT!!!!!!!!!!**
5. ADDITIONAL INSTRUCTIONS: _____

DAY OF EXAMINATION

1. You may have clear liquids until 4 hours prior to your procedure. You may take your regular medications with a sip of water up until your procedure time.
2. You should plan on spending up to 2 hours at our facility for this test.

Thank you for the opportunity to accommodate your healthcare needs.

(03/18)